

Name: _____ Date of Hatching: _____ Qty: _____ chicks

Vaccination guideline for Indigenous chicken

Age	Vaccine	Mode of administration	Remarks
Day old	Mareks	<ul style="list-style-type: none"> • Subcutaneous 	Mainly for commercial hatcheries Recommended Vaccine: NOBILIS RISMAVAC
Day 14	Gumboro (1 st Dose)	<ul style="list-style-type: none"> • Drinking water 	Recommended Vaccine: HIPRAGUMBORO-CH/80
3 weeks (Day 21)	Newcastle disease (1 st Dose)	<ul style="list-style-type: none"> • Eye drop or • Drinking water 	(Repeat after two months) Recommended vaccine: HIPRAVIAR B1 / V.H. + H-120
Day 24	Gumboro (2 nd Dose)	<ul style="list-style-type: none"> • Drinking water 	Recommended Vaccine: HIPRAGUMBORO-CH/80
3 weeks (In hot areas)	Fowl pox	<ul style="list-style-type: none"> • Wing stab 	Recommended vaccine: HIPRAPOX

6 weeks (other areas)	Fowl pox	<ul style="list-style-type: none"> • Wing stab 	Recommended vaccine: HIPRAPOX
8 weeks	Newcastle Disease (2 nd Dose)	<ul style="list-style-type: none"> • Eye drop or • Drinking water 	Recommended vaccine: HIPRAVIAR B1 / V.H. + H-120
	Fowl typhoid	<ul style="list-style-type: none"> • Intramuscular injection 	FowlVax by KEVIVAPI(Kenya Veterinary Vaccines Production Institute)
18 weeks	Newcastle Disease (3 rd Dose)	<ul style="list-style-type: none"> • Eye drop or • Drinking water 	Recommended vaccine: HIPRAVIAR B1 / V.H. + H-120 Repeat every 3 months
19 weeks	De-worming	<ul style="list-style-type: none"> • Drinking water 	Recommended: Levacide Poultry Repeat every 3 months

Notes.

- **NEVER** vaccinate sick chicken
- Consult your veterinary/livestock staff for detailed vaccination programs in your area

HOMERANGE